

# PRV

PATENT- OCH REGISTRERINGSVERKET

Patentavdelningen

REC'D 15 APR 2005

WIPO

PCT

IB/05/00673

## Intyg Certificate

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.



(71) Sökande                      Pfizer Health AB, Stockholm SE  
Applicant (s)

(21) Patentansökningsnummer    0400685-4 ✓  
Patent application number

(86) Ingivningsdatum                      2004-03-19 ✓  
Date of filing

Stockholm, 2005-01-19

För Patent- och registreringsverket  
For the Patent- and Registration Office

Hjärdís Segerlund

Avgift

Fee                      170:-

## PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

PATENT- OCH  
REGISTRERINGSVERKET  
SWEDEN

Postadress/Adress  
Box 5055  
S-102 42 STOCKHOLM

Telefon/Phone  
+46 8 782 25 00  
Vx 08-782 25 00

Telex  
17978  
PATOREG S

Telefax  
+46 8 666 02 86  
08-666 02 86

## MEANS FOR TRANSDERMAL ADMINISTRATION OF NICOTINE

### Technical Field

The present invention relates to devices for transdermal administration of nicotine, specifically such devices providing for basic as well as additional user activatable administration of nicotine. The present invention also relates to use of said devices.

### Background of the Invention

#### *Tobacco dependence and reduction thereof*

In recent years, with the recognition of the harmful effects of tobacco smoking, there have been numerous campaigns and programs by governmental agencies and various health groups and other interested organisations to disseminate information about the adverse health effects resulting from tobacco smoking. Moreover, and as a result of this recognition of the harmful effects, there have been many programs directed to attempts in reducing smoking incidence.

Nicotine is an organic compound and is the principal alkaloid of tobacco. Nicotine is the chief addictive ingredient in the tobacco used in cigarettes, cigars, snuff and the like. Nicotine is also an addictive drug, though, and smokers characteristically display a strong tendency to relapse after having successfully stopped smoking for a time. Nicotine is the worlds second most used drug, after caffeine from coffee and tea.

The main problem with tobacco smoking is its enormous implications on health. It is estimated that smoking related diseases cause some 3 – 4 million deaths per year globally. According to Centers for Disease Control and Prevention. Cigarette smoking among adults – United States, 1995. MMWR 1997;46:1217 – 1220 around 500,000 persons in USA die each year as a result of tobacco use. In fact, excessive smoking is now recognised as one of the major health problems throughout the world. This grim consequence of tobacco smoking has urged many medical associations and health authorities to take very strong actions against the use of tobacco.

Even though tobacco smoking is decreasing in many developed countries today it is hard to see how the societies could get rid of the world's second most used drug.

The most advantageous thing a heavy smoker can do is to reduce or preferably even stop smoking completely. Experience shows, however, that most smokers find this extremely difficult since, mostly, tobacco smoking results in a dependence disorder or craving. The WHO has in its International Classification of Disorders a diagnosis called Tobacco

Dependence. Others like the American Psychiatric Association call the addiction Nicotine Dependence. It is generally accepted that these difficulties to stop smoking result from the fact those heavy smokers are dependent on nicotine. The most important risk factors are, however, substances that are formed during the combustion of tobacco, such as carbon  
 5 monoxide, tar products, aldehydes, and hydro cyanic acid.

#### *Effects of nicotine*

The administration of nicotine can give satisfaction and the usual method is by smoking, either by smoking e g a cigarette, a cigar or a pipe. However, smoking has health hazards and it is therefore desirable to formulate an alternative way of administering nico-  
 10 tine in a pleasurable manner that can be used to facilitate withdrawal from smoking and/or used as a replacement for smoking.

When smoking a cigarette, nicotine is quickly absorbed into the smoker's blood and reaches the brain within around ten seconds after inhalation. The quick uptake of nicotine gives the consumer a rapid satisfaction, or kick. The satisfaction, then, lasts during the  
 15 smoking time of the cigarette and for a period of time thereafter. The poisonous, toxic, carcinogenic, and addictive nature of smoking has provided efforts for methods, compositions and devices, which help in breaking the habit of smoking cigarettes.

Nicotine is an addictive poisonous alkaloid  $C_5H_4NC_4H_7NCH_3$ , derived from the tobacco plant. Nicotine is also used as an insecticide.

#### *Nicotine replacement products*

One way to reduce smoking is to provide nicotine in a form or manner other than by smoking and some products have been developed to fulfil this need. Nicotine containing formulations are currently the dominating treatments for tobacco dependence.

The successes in achieving reduction in the incidence of smoking have been relatively poor using presently known products. The present state of the art involves both behavioural approaches and pharmacological approaches. More than 80 % of the tobacco  
 25 smokers who initially quit smoking after using some behavioural or pharmacological approach to singly reduce smoking incidence generally relapse and return to the habit of smoking at their former rate of smoking within about a one year's period of time.

30 As an aid for those who are willing to stop smoking there are several ways and forms of nicotine replacement products available on the market, such as nicotine chewing gum. Several methods and means have been described for diminishing the desire of a subject to use tobacco, which comprises the step of administering to the subject nicotine or a de-

rivative thereof as described in e g US 5,810,018 (oral nicotine spray), US 5,939,100 (nicotine containing microspheres) and US 4,967,773 (nicotine containing lozenge).

Nicotine-containing nose drops have been reported (Russell et al., British Medical Journal, Vol. 286, p. 683 (1983); Jarvis et al., Brit. J. of Addiction, Vol. 82, p. 983 (1987)).

- 5 Nose drops, however, are difficult to administer and are not convenient for use at work or in other public situations. Ways of administering nicotine by way of delivering directly into the nasal cavity by spraying is known from US 4,579,858, DE 32 41 437 and WO93/127 64. There may, though, be local nasal irritation with use of nasal nicotine formulations. The difficulty in administration also results in unpredictability of the dose of nicotine admini-  
10 stered.

Also, inhaling devices resembling a cigarette are known for uptake of nicotine vapours as suggested in US 5,167,242.

One successful approach to date in reducing the incidence of smoking relies upon nicotine containing chewing gum that is designed to reduce smoking withdrawal symptoms.

- 15 The reported success rate is approximately twice that of placebo.

One successful product that is used as a smoking substitute and/or as a smoking cessation aid and which is based on nicotine, is the chewing gum Nicorette®. This product was one of the first nicotine replacement forms that was approved by the Food and Drug Administration (FDA) and is still one of the most used nicotine replacement products.

- 20 Nicorette® chewing gum has been on the market in about 60 countries for several years. In this chewing gum the nicotine is present in the form of a complex with an insoluble cation-exchanger (polacrilex) that is dispersed in a gum base. The nicotine is slowly released from the gum due to chewing and will reach similar plasma levels as when smoking a cigarette after about 30 minutes depending on the chewing technique, i e slow or active chewing.

- 25 Patents related to this product are e g US Patents 3,877,468, 3,901,248 and 3,845,217.

#### *Nicotine transdermal patches*

The use of skin patches for transdermal administration of nicotine was reported many years ago (Rose, in Pharmacologic Treatment of Tobacco Dependence, (1986) pp.

- 158-166, Harvard Univ. Press). A large number of patents on devices for transdermal delivery of nicotine have been issued, e g US 5,120,546 disclosing a transdermal delivery system wherein the nicotine is complexed to a cyclo compound, US 5,230,896 disclosing a transdermal delivery system wherein is made use of an acrylic polymer adhesive, US 4,943,435 disclosing a transdermal patch for delivering nicotine for 12 – 24 hours making use of a  
30

rate-controlling membrane, US 4,915,950 disclosing a method for making nicotine-containing transdermal delivery devices whereby the nicotine is printed onto an adsorbent fabric layer, and US 5,533,995 disclosing a transdermal device wherein the nicotine transport within the device as such is controllable using an internal electrode system. The aforesaid  
 5 patents relate to transdermal devices wherein the drug flux through the skin cannot as such be regulated by the user.

*Prior art and problems thereof*

There are known methods for providing an essentially constant base-line plasma level of nicotine being supplemented with boost doses of nicotine when the patient so de-  
 10 sires. These methods to date imply use of two different dosage forms for delivery of nicotine to be used in combination. For this purpose may e g be used a nicotine patch in combination with a nicotine lozenge, see US 5,135,753, or a nicotine patch in combination with a nicotine nasal spray, see US 5,721,257. For convenience of the user there is a need to provide a single dosage form, which may provide an essentially constant base-line plasma level  
 15 of nicotine as well as additional boost doses of nicotine when the patient/user so desires.

The present invention pertains to a solution to the captioned problem by providing a unitary device, which accounts for a basic as well as an additional user activatable transdermal delivery of nicotine.

It is highly desirable in light of the aforementioned problems to develop means and  
 20 methods for the administration of nicotine to provide satisfaction to a person craving for nicotine or to provide a sense of smoking satisfaction without smoking, which can also avoid problems associated with the prior art means and methods. In this respect, the present invention addresses this need and interest.

**Definitions**

25 With "basic transdermal administration of nicotine in any form" and similar expressions is herein meant transdermal nicotine transport provided by a part of the claimed device that administers nicotine continuously during the period of the intended application of the claimed device on the skin.

30 With "additional transdermal administration of nicotine in any form" and similar expressions is herein meant transdermal nicotine transport provided by a part of the claimed device that at user activation administers additional nicotine, beyond that provided by the part of the claimed device providing for basic administration of nicotine in any form.

### **Legend of Figures**

- Figure 1 is a schematic drawing of the embodiment according to Example 1.  
 Figure 2 is a schematic drawing of the embodiment according to Example 2.  
 Figure 3 is a schematic drawing of the embodiment according to Example 3.  
 5 Figure 4 is a schematic drawing of the embodiment according to Example 4.  
 Figure 5 is a schematic drawing of the embodiment according to Example 5.  
 Figure 6 is a schematic drawing of the embodiment according to Example 6.

### **Summary of the Invention**

10 In view of the foregoing disadvantages known in the art when trying to deliver nicotine to a subject to provide different delivery rates of nicotine to a subject the present invention provides novel devices for combined basic and additional user activatable transdermal delivery of nicotine.

Further, the present invention provides use of nicotine for the manufacture of a device, and a method for aiding in smoking cessation, in temporary smoking abstinence  
 15 and/or in reducing the urge to smoke or to otherwise use tobacco containing material, and/or for treating conditions suitable for treatment with nicotine, such conditions being selected from the group consisting of Alzheimer's disease, Crohn's disease, Parkinson's disease, Tourette's syndrome, ulcerous colitis and post-smoking-cessation weight control by transdermal administration of nicotine with such a device. The method of transdermal  
 20 delivery of nicotine can also be practiced in combination with means for delivering of nicotine selected from mouth sprays, nasal sprays, transdermal patches, inhaling devices, lozenges, tablets and from parenteral methods, subcutaneous methods, intravenous methods, rectal methods, vaginal methods and transmucousal methods, including use of tobacco.

### **25 Categorization of transdermal devices**

Means for transdermal administration of nicotine in any form can be categorized in many different ways. A comprehensive categorization of transdermal devices useful in the present invention is in four main groups as follows:

- the reservoir type, in which the drug is placed in a liquid or a gel and delivered  
 30 across a rate-moderating membrane to the skin;
- the matrix type, in which the drug is placed within a non-adhesive polymeric material, typically a hydrogel or soft polymer;
- the drug-in-adhesive type, in which the drug is placed within an adhesive polymer;

- the multi-laminate type, which is similar to the drug-in-adhesive design, but which incorporates an additional layer of pressure sensitive adhesive to cover the entire device and affix it to the skin. A membrane can also be incorporated into the multi-laminate type.

**User activatable delivery of nicotine**

5 Means for modifying transdermal flux of pharmaceutically active substances can be categorized in many different ways. One such categorization is according to below Table 2.

Since many of the means listed in Table 2 may be designed to be both activated and deactivated by the user, they may be well suited to be used for additional transdermal administration of nicotine in the present invention. Means especially suited for user activation  
10 and deactivation, and thus preferred for use in additional transdermal administration of nicotine of the present invention include, but are not limited to: iontophoresis (including also electromigration and electroosmosis), sonophoresis, micro-needles, jet injection or combinations thereof.

**Table 2. Means for modifying transdermal delivery**

15

Means	Mechanism
Iontophoresis (electromigration, electroosmosis)	Transport of charged drug molecules in an electrical field drives drug molecules through the Stratum Corneum (SC), enhancing transdermal transport rate.
Electroporation	Transport-enhancing pores through SC created by an electrical current.
Sonophoresis	Transport enhancement by heating and/or disordering SC by means of ultrasound.
Micro-needles	Microscopic needles, loaded with drug, punctures the SC thus increasing transdermal drug flux.
Chemical enhancers	Chemicals interacting with the structure of SC, leading to increased permeability of drug molecules through SC.
Carrier particles (e.g. transfersomes, lipopearls)	Drug enveloped in submicron particles with good SC permeability.
Jet injection	Rapidly expanding gas "shoots" drug particles through SC, leading to increased permeability of drug molecules.
Laser microporation	Laser light punctures SC, leading to increased perme-

	ability of drug molecules.
Tape stripping	Adhesive tape strips off SC, leading to increased permeability of drug molecules.
Suction ablation	Suction device strips off SC, leading to increased permeability of drug molecules.
Metabolic inhibitors	Chemicals that slows the reparative processes of the skin. Damaged SC more permeable to drug than undamaged SC.
Supersaturation	Maximizes the concentration gradient at the drug/SC interface thus optimizing transdermal diffusion of drug.
Occlusion	Transport enhancement through heating and/or disordering SC by means of an occlusive dressing.
Solvent drag	The transdermal flux of a highly permeable substance drags drug molecules along with it, thus increasing the transdermal transport rate of the drug.

The means of Table 2 are as such known in the art.

#### **The active ingredient**

According to the invention, the claimed device comprises nicotine in any form.

- 5 Nicotine is intended to include nicotine, 3-(1-methyl-2-pyrrolidinyl)-pyridine, with its base form, including synthetic nicotine as well as nicotine extracts from tobacco plants, or parts thereof, such as the genus *Nicotiana* alone or in combination.

- Nicotine in any form is selected from the group consisting of a nicotine salt, the free base form of nicotine, a nicotine derivative, such as a nicotine cation exchanger, a nicotine inclusion complex or nicotine in any non-covalent binding; nicotine bound to zeolites; nicotine bound to cellulose or nicotine bound to starch microspheres; and mixtures thereof.
- 10

Numerous nicotine salts are known, and may be used, e g the salts presented in below Table 3.

**Table 3 Acids useful for nicotine salt formation**

Acid	Molar ratio* of acid:nicotine
Formic	2:1



Acetic	3:1
Propionic	3:1
Butyric	3:1
2-Methylbutyric	3:1
3-Methylbutyric	3:1
Valeric	3:1
Lauric	3:1
Palmitic	3:1
Tartaric	2:1
Citric	2:1
Malic	2:1
Oxalic	2:1
Benzoic	1:1
Gentisic	1:1
Gallic	1:1
Phenylacetic	3:1
Salicylic	1:1
Phthalic	1:1
Picric	2:1
Sulfosalicylic	1:1
Tannic	1:5
Pectic	1:3
Alginic	1:2
Hydrochloric	2:1
Chloroplatinic	1:1
Silicotungstic	1:1
Pyruvic	2:1
Glutamic	1:1
Aspartic	1:1

---

\* recommended upon production

The above-mentioned cation exchanger is preferably a polyacrylate.

The above-mentioned inclusion complex is preferably a complex with a cyclodextrin, such as  $\beta$ -cyclodextrin.

The above-mentioned nicotine salt is preferably a tartrate, hydrogen tartrate, citrate, maleate or hydrochloride.

- 5        The form of nicotine being provided through the basic administration may be another form of nicotine than the one being provided through the additional administration.

**Further additives**

Further additives may be added optionally to a device according to the present invention.

- 10        Optional additives preferably comprise one or more additives selected from the group consisting of stabilisers, enhancers and anti-irritants.

Stabilizers may be selected from the group consisting of antioxidants including vitamin E, i.e. tocopherole, ascorbic acid, sodium pyrosulfite, or butylated hydroxytoluene (BHT), butylated hydroxyanisole, edetic acid and edetate salts; and preservatives including

- 15        citric acid, tartaric acid, lactic acid, malic acid, acetic acid, benzoic acid, and sorbic acid. Preferred embodiments comprise an antioxidant as the stabiliser, and even more preferably vitamin E and/or butylated hydroxytoluene (BHT).

Enhancers may be selected from the group consisting of

- alcohols, such as short chain alcohols, e.g. ethanol and the like, long chain fatty alcohols,
- 20        e.g. lauryl alcohols, and the like, and polyalcohols, e.g. propylene glycol, glycerin and the like;
- amides, such as amides with long aliphatic chains, or aromatic amides like N,N-diethyl-m-toluamide;
- amino acids;
- 25        - azone and azone-like compounds;
- essential oils, i.e. essential oils or constituents thereof, such as l-carvone, l-menthone and the like;
- fatty acids and fatty acid esters, such as oleic acid, lauric acid and the like, further esters of fatty acids, such as isopropyl myristate, and various esters of lauric acid and of oleic acid
- 30        and the like;
- macrocyclic compounds, such as cyclopentadecanone and cyclodextrins;
- phospholipid and phosphate compounds, such as phospholipids;
- 2-pyrrolidone compounds; and

- miscellaneous compounds, like sulfoxides, such as dimethyl sulfoxides, and fatty acid ethers, such as Laureth-9 and polyoxylaurylether.

Combinations of enhancers from different groups may prove useful and efficient.

An example of a useful anti-irritant is vitamin E.

## 5 Examples

The below examples are illustrative and non-limiting. The numbers within brackets refer to the corresponding numbers in the figures.

Example 1, being schematically illustrated in Figure 1.

## 10 Basic administration from matrix part and additional administration from iontophoretic part.

A device in the format of a transdermal patch consisting of two different nicotine delivery systems:

One system where the transdermal dose of nicotine is delivered from a polymeric matrix compartment (11). The mechanism for transdermal nicotine delivery for this system is passive diffusion along a concentration gradient. The rate of delivery is governed by the properties of the matrix polymers and the concentration of nicotine loaded into the matrix. The nicotine in the matrix is in its neutral, base form.

One system where the transdermal dose of nicotine is delivered from a compartment where the mechanism of nicotine transport is iontophoresis, i.e. the driving force is supplied by an electric current (12,13). In this system positively charged nicotine is loaded in the electrode-fitted (anode) drug compartment (12). The system is completed by a second electrode-fitted (cathode) compartment (13) where negatively charged counter-ions, preferably chloride ions but also other biocompatible negatively charged ions may be used, are loaded. Electrodes (14) are preferably of the Ag/AgCl type. A disposable battery (14), capable of outputting a current of up to 0.5 mA/cm<sup>2</sup> of drug compartment area in contact with the skin, is connected to the electrodes. When the iontophoretic system is activated, the skin tissue at the application site completes the electrical circuit. The rate of nicotine delivery for this system is governed by the size of the electrical current between the electrodes (14).

The two above systems are placed side-by-side in a thin patch-like device 20-50 cm<sup>2</sup> in size. The device is fastened at the application site by means of an adhesive layer covering the bottom of the device. The device is backed by a flexible non-woven material, which provides structural support. Directly below the non-woven material is a thin aluminum

layer, acting as a barrier to nicotine diffusion through the backing material of the device. The device is also fitted with an activation button (15) to initiate the driving electrical current of the iontophoretic system. The passive diffusion system provides a basic dose of nicotine. The user may activate the iontophoretic system by pressing the activation button (15), and thus release an additional dose of nicotine. Deactivation of the iontophoretic system is by a timer function. Alternatively, the user may at any time deactivate the additional administration of nicotine by pressing the activation button (15) a second time.

Example 2, being schematically illustrated in Figure 2.

Basic administration from reservoir part and additional administration from micro-needle part.

A device in the format of a transdermal patch consisting of two different nicotine delivery systems:

One system where the transdermal dose of nicotine is delivered from a reservoir compartment, through a rate-controlling membrane (21). The mechanism for transdermal nicotine delivery for this system is passive diffusion along a concentration gradient. The rate of delivery is governed by the properties of the rate-controlling membrane and the concentration of nicotine loaded into the compartment. The nicotine in the matrix is in its neutral, base form.

One system where the transdermal dose of nicotine is delivered from a reservoir compartment lined on one side with micro-needles with the function of being able to puncture the stratum corneum of the skin, thus increasing the transdermal delivery rate of nicotine (22). The micro-needles are suspended a short distance from the skin surface and the user may activate the system by pressing on the backing of the patch to bring the micro-needles in contact with the skin. The rate of delivery is governed by the properties of the micro-needles and the concentration of nicotine loaded into the compartment. The nicotine in the micro-needle reservoir compartment is in its neutral, base form.

The two above systems are placed side-by-side in a thin patch-like device 20-50 cm<sup>2</sup> in size. The device is backed by a flexible non-woven material, which provides structural support. Directly below the non-woven material is a thin aluminum layer, acting as a barrier to nicotine diffusion through the backing material of the device. The device is fastened at the application site by means of an adhesive layer covering the bottom of the device. The immediate area surrounding the micro-needle part of the device consists of a semi-rigid flexing polymer (23) that will allow the micro-needles to flex in and out of contact with the

stratum corneum (24) dictated by the user pressing on the backing of the device, directly behind the micro-needles. The passive diffusion system provides a basic dose of nicotine. The user may activate an additional dose of nicotine by pressing on the device backing so that the micro-needles puncture the stratum corneum (24) of the skin and channel nicotine

5 through the stratum corneum (24) to the underlying tissue.

Example 3, being schematically illustrated in Figure 3.

Basic administration from matrix part and additional administration from ultrasonic part.

10 A device in the format of a transdermal patch consisting of two different nicotine delivery systems:

One system where the transdermal dose of nicotine is delivered from a polymeric matrix compartment (31). The mechanism for transdermal nicotine delivery for this system is passive diffusion along a concentration gradient. The rate of delivery is governed by the properties of the matrix polymers and the concentration of nicotine loaded into the matrix.

15 The nicotine in the matrix is in its neutral, base form.

One system where the transdermal dose of nicotine is delivered from a polymeric matrix compartment (32) where the mechanism of nicotine transport is sonophoresis, i.e. an ultrasonic device (33) causes heating of the skin and a reversible disordering of the structure of the stratum corneum. The rate of nicotine delivery for this system is governed by the

20 intensity of the ultrasonic vibrations. The nicotine in the matrix is in its neutral, base form.

The two above systems are placed side-by-side in a thin patch-like device 20-50 cm<sup>2</sup> in size. The device is fastened at the application site by means of an adhesive layer covering the bottom of the device. The device is also fitted with an activation button (34) to initiate the ultrasonic vibrations. The passive diffusion system provides a basic dose of nicotine. The

25 user may activate the sonophoretic system by pressing the activation button (34), and thus release an additional dose of nicotine. Deactivation of the sonophoretic system is by a timer function. Alternatively, the user may at any time deactivate the additional administration of nicotine by pressing the activation button (34) a second time.

Example 4, being schematically illustrated in Figure 4.

30 Basic administration from matrix part sharing space with iontophoretic part providing additional administration.

A device in the format of a transdermal patch consisting of two different nicotine delivery systems:

One system where the transdermal dose of nicotine is delivered from a compartment where the mechanism of nicotine transport is iontophoresis, i.e. the driving force is supplied by an electric current. In this system positively charged nicotine is loaded in the electrode-fitted (anode) drug compartment (41). The system is completed by a second electrode-fitted (cathode) compartment (42) where negatively charged counter-ions are loaded. This compartment also contains the neutral base nicotine of the passive diffusion system detailed below. A battery (43) supplies the needed electrical power. The rate of nicotine delivery for this system is governed by the size of the electrical current between the electrodes.

One system where the transdermal dose of nicotine is delivered from a polymeric matrix compartment (42). This compartment contains both nicotine in its neutral, base form and the negatively charged counter-ions needed for the iontophoretic system. The mechanism for transdermal nicotine delivery for this system is passive diffusion along a concentration gradient. The rate of nicotine delivery is governed by the properties of the matrix polymers and the concentration of nicotine loaded into the matrix.

The two above systems are fitted in a thin patch-like device 20-50 cm<sup>2</sup> in size. The device is fastened at the application site by means of an adhesive layer covering the bottom of the device. The device is backed by a flexible non-woven material, which provides structural support. Directly below the non-woven material is a thin aluminum layer, acting as a barrier to nicotine diffusion through the backing material of the device. The device is also fitted with an activation button (44) to initiate the driving electrical current of the iontophoretic system. The passive diffusion part, located in the iontophoretic cathode compartment (42), provides a basic dose of nicotine. The user may activate the iontophoretic system by pressing the activation button (44), and thus release an additional dose of positively charged nicotine from the iontophoretic anode compartment (41). Deactivation of the iontophoretic system is by a timer function. Alternatively, the user may at any time deactivate the additional administration of nicotine by pressing the activation button (44) a second time.

Example 5, being schematically illustrated in Figure 5.

Basic administration from reservoir part and additional administration from iontophoretic part, where the two parts may be individually applied to the skin.

A device in the format of a transdermal patch consisting of two different nicotine delivery systems:

One system where the transdermal dose of nicotine is delivered from a reservoir compartment, through a rate-controlling membrane (51). The mechanism for transdermal nicotine delivery for this system is passive diffusion along a concentration gradient. The rate of delivery is governed by the properties of the rate-controlling membrane and the concentration of nicotine loaded into the compartment. The nicotine in the matrix is in its neutral, base form.

One system where the transdermal dose of nicotine is delivered from a compartment where the mechanism of nicotine transport is iontophoresis, i.e. the driving force is supplied by an electric current. In this system positively charged nicotine is loaded in an electrode-fitted drug compartment (52). The system is completed by a second electrode-fitted compartment where negatively charged counter-ions are loaded (53). A battery (54) supplies the needed electrical power. The rate of nicotine delivery for this system is governed by the size of the electrical current between the electrodes.

The two above systems are placed side-by-side in a thin patch-like device 20-50 cm<sup>2</sup> in size. The device is fastened at the application site by means of an adhesive layer covering the bottom of the device. The device is backed by a flexible non-woven material, which provides structural support. Directly below the non-woven material is a thin aluminum layer, acting as a barrier to nicotine diffusion through the backing material of the device. The two systems are separated by a perforation (55) in the device, making it possible to tear off one system from the other for individual placement on the body of the two parts. The iontophoretic part of the device is also fitted with an activation button (56) to initiate the driving electrical current of the iontophoretic system. The passive diffusion system provides a basic dose of nicotine. The user may activate the iontophoretic system by pressing the activation button (56), and thus release an additional dose of nicotine. Deactivation of the iontophoretic system is by a timer function. Alternatively, the user may at any time deactivate the additional administration of nicotine by pressing the activation button (56) a second time.

Example 6, being schematically illustrated in Figure 6.

Basic administration from matrix part and additional, multilevel administration from iontophoretic part.

A device in the format of a transdermal patch consisting of two different nicotine delivery systems:

One system where the transdermal dose of nicotine is delivered from a polymeric matrix compartment (61). The mechanism for transdermal nicotine delivery for this system is passive diffusion along a concentration gradient. The rate of delivery is governed by the properties of the matrix polymers and the concentration of nicotine loaded into the matrix.

5 The nicotine in the matrix is in its neutral, base form.

One system where the transdermal dose of nicotine is delivered from a compartment where the mechanism of nicotine transport is iontophoresis, i.e. the driving force is supplied by an electric current. In this system positively charged nicotine is loaded in the electrode-fitted (anode) drug compartment (62). The system is completed by a second electrode-fitted  
10 (cathode) compartment (63) where negatively charged counter-ions are loaded. A battery (64) supplies the needed electrical power. The rate of nicotine delivery for this system is governed by the size of the electrical current between the electrodes.

The two above systems are placed side-by-side in a thin patch-like device 20-50 cm<sup>2</sup> in size. The device is fastened at the application site by means of an adhesive layer covering  
15 the bottom of the device. The device is backed by a flexible non-woven material, which provides structural support. Directly below the non-woven material is a thin aluminum layer, acting as a barrier to nicotine diffusion through the backing material of the device. The passive diffusion system provides a basic dose of nicotine. The device is also fitted with a set of activation buttons (65,66,67) to initiate the driving electrical current of the iontophoretic system. The user may activate the iontophoretic system by pressing one of the activa-  
20 tion buttons (65,66,67), and thus release an additional dose of nicotine. The amount of nicotine of the additional dose ranges from low to high, depending on which button is pressed by the user. Each button corresponds to a pre-set dose level. Deactivation of the iontophoretic system is by a timer function. Alternatively, the user may at any time deac-  
25 tivate the additional administration of nicotine by pressing the same activation button a second time.

#### **Further embodiments**

Other combinations of herein disclosed means for obtaining the basic admini-  
stration and the additional administration respectively than those disclosed in the Examples  
30 are readily envisagable and useful.

The present device should preferably be occlusive.

The present device may deliver nicotine during a predefined period of time, preferably 12, 16, 24 or 48 hours.



The at least one part providing for basic transdermal administration and the at least one part providing for additional and user activatable transdermal administration may be combined in different ways. For example may said parts have at least one feature in common, e g a common adhesive layer and/or a common drug reservoir. Said parts may also be  
5 combined so that they have no features in common.

# CLAIMS

1. Device for transdermal administration of nicotine in any form, characterized in that it comprises at least one first part providing for basic administration of nicotine in any form and at least one second part providing for additional administration of nicotine in any form, the at least one second part being activatable by the user.
2. Device for transdermal administration according to claim 1, characterized in that the at least one first part providing for basic administration is/are (a) transdermal patch(es), preferably of the reservoir type, the matrix type, the drug-in-adhesive type and/or the multi-laminate type, preferably of the drug-in-adhesive type or the reservoir type or combinations of these two types.
3. Device for transdermal administration according to any preceding claim, characterized in that the at least one second part providing for additional administration comprise(s) means for iontophoretic delivery, sonophoresis, jet injection and/or micro-needles.
4. Device for transdermal administration according to claim 3, characterized in that the at least one second part providing for additional administration comprise(s) means for iontophoretic delivery.
5. Device for transdermal administration according to any preceding claim, characterized in that the at least one first part and the at least one second part have at least one feature in common.
6. Device for transdermal administration according to any of claims 1 - 5, characterized in that the at least one first part and the at least one second part are detachable from one another and/or may be individually applied.
7. Device for transdermal administration according to any preceding claim, characterized in that the form of nicotine is selected from nicotine free base, a nicotine salt, such as a tartrate, hydrogen tartrate, citrate, maleate or hydrochloride, a nicotine inclusion complex, such as a complex with a cyclodextrin, a nicotine cation exchanger, such as nicotine with polyacrylate or nicotine in any non-covalent binding, nicotine bound to zeolites, nicotine bound to cellulose or nicotine bound to starch microspheres, and mixtures thereof.
8. Device for transdermal administration according to claim 7, characterized in that the form of nicotine is nicotine free base.

9. Device for transdermal administration according to claim 7 or 8, characterized in that the least one part providing for basic administration of nicotine is for delivery of nicotine in another form than the at least one part providing for additional administration of nicotine

5 10. Device for transdermal administration according to anyone of the preceding claims, characterized in that it delivers nicotine during a predefined period of time, preferably 12, 16, 24 or 48 hours.

11. Device according to anyone of the preceding claims, characterized in that it further comprises one or more stabilisers, preferably selected from the group consisting of antioxidants including vitamin E, i.e. tocopherole, ascorbic acid, sodium pyrosulfite, butylated hydroxytoluene (BHT), butylated hydroxyanisole, edetic acid and edetate salts, and preservatives including citric acid, tartaric acid, lactic acid, malic acid, acetic acid, benzoic acid, and sorbic acid, preferably vitamin E and/or butylated hydroxytoluene (BHT).

12. Device according to anyone of the preceding claims, characterized in that it further comprises one or more substances enhancing transdermal penetration, preferably a compound selected from the group consisting of

- alcohols, such as short chain alcohols, e.g. ethanol and the like, long chain fatty alcohols, e.g. lauryl alcohols, and the like, and polyalcohols, e.g. propylene glycol, glycerin;
- amides, such as amides with long aliphatic chains, or aromatic amides like N,N-diethyl-m-toluidamide;
- amino acids;
- azone and azone-like compounds;
- essential oils, i.e. essential oils or constituents thereof, such as l-carvone, l-menthone and the like;
- fatty acids and fatty acid esters, such as oleic acid, lauric acid and the like, further esters of fatty acids, such as isopropyl myristate, and various esters of lauric acid and of oleic acid;
- macrocyclic compounds, such as cyclopentadecanone and cyclodextrins;
- phospholipid and phosphate compounds, such as phospholipids;
- 2-pyrrolidone compounds; and
- miscellaneous compounds, like sulphoxides, such as dimethyl sulphoxides, and fatty acid ethers, such as Laureth-9 and polyoxy-laurylether.

13. Device according to anyone of the preceding claims, characterized in that it further comprises one or more substances reducing irritant reactions, preferably vitamin E.

14. Device according to anyone of the preceding claims, characterized in  
5 that it is occlusive.

15. Use of nicotine for the manufacture of a device for aiding in smoking cessation, in temporary smoking abstinence and/or in reducing the urge to smoke or to otherwise use tobacco containing material, and/or for treating conditions suitable for treatment with nicotine, such conditions being selected from the group consisting of Alzheimer's disease,  
10 Crohn's disease, Parkinson's disease, Tourette's syndrome, ulcerous colitis and post-smoking-cessation weight control, such a device being according to anyone of claims 1 - 14.

16. Method for aiding in smoking cessation, in temporary smoking abstinence and/or in reducing the urge to smoke or to otherwise use tobacco containing material,  
15 and/or for treating conditions suitable for treatment with nicotine, such conditions being selected from the group consisting of Alzheimer's disease, Crohn's disease, Parkinson's disease, Tourette's syndrome, ulcerous colitis and post-smoking-cessation weight control, by transdermal administration of nicotine with a device according to anyone of claims 1 - 14.

17. Method for aiding in smoking cessation, in temporary smoking abstinence and/or in reducing the urge to smoke or to otherwise use tobacco containing material, and/or for treating conditions suitable for treatment with nicotine, such conditions being selected from the group consisting of Alzheimer's disease, Crohn's disease, Parkinson's disease, Tourette's syndrome, ulcerous colitis and post-smoking-cessation weight control,  
20 by transdermal administration of nicotine with a device according to anyone of claims 1 - 14 in combination with means selected from mouth sprays, nasal sprays, transdermal patches, inhaling devices, lozenges, tablets and from parenteral methods, subcutaneous methods, intravenous methods, rectal methods, vaginal methods and transmucousal methods, including use of tobacco.


**ABSTRACT**

Devices for transdermal administration of nicotine, which provide for basic as well as additional user activatable administration of nicotine, as well as use of such devices.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
221


1/6

Figure 1




2/6

Figure 2



3/6


Figure 3






4/6

Figure 4





6/6

Figure 6

